

Platelet Activation Induced by Porcine Factor VIII (HYATE:C)

H. Chang,¹ M. Mody,¹ A.H. Lazarus,¹ F. Ofofu,³ M.B. Garvey,¹ V. Blanchette,² J. Teitel,¹ and J. Freedman^{1*}

¹St. Michael's Hospital University of Toronto, Toronto, Canada

²Hospital for Sick Children, University of Toronto, Toronto, Canada

³McMaster University Medical Center, Hamilton, Ontario, Canada

We studied the effects of porcine factor VIII (P-FVIII; Hyate:C) and other coagulation products employed in the management of patients with hemophilia A, on platelet activation *in vitro*. Exposure of normal resting platelets to P-FVIII resulted in platelet activation, as manifested by increased expression of the platelet surface activation markers CD62, CD63, and activated-GPIIb/IIIa, and by activation-induced modulation of expression of normal platelet membrane glycoproteins CD41, CD42, and CD36. In contrast, platelet activation was not observed after exposure of the platelets to human FVIII, FEIBA, recombinant FVIIIa, or cryosupernatant plasma. As with thrombin, exposure of platelets to P-FVIII resulted in the generation of platelet microparticles, an effect not seen with the other products. In contrast to the characteristic reduction in expression in the number of CD42 molecules detected on thrombin-activated platelets, P-FVIII-stimulated platelets showed a small increase in CD42 expression. In contrast to thrombin, P-FVIII did not cause platelet dense granule release. The results indicate that therapeutic P-FVIII activates platelets, likely in ways that are different from the platelet activation seen with thrombin. The observed platelet activation and microparticle generation may provide a "hypercoagulable" mechanism for hemostasis with P-FVIII therapy separate from, and additional to, that due to increased circulating FVIII levels. *Am. J. Hematol.* 57:200–205, 1998. © 1998 Wiley-Liss, Inc.

Key words: platelets; platelet activation; porcine factor VIII; hemophilia

INTRODUCTION

Porcine factor VIII (P-FVIII; Hyate:C) concentrates are used in the management of hemophilic patients with inhibitors to human factor VIII. Early preparations of P-FVIII were associated with a decrease in platelet count post-infusion [1,2]. The thrombocytopenia has been attributed to platelet aggregation, caused by porcine von Willebrand Factor (vWF) present in the P-FVIII product on human platelets [1,3–5]. With more purified P-FVIII products, a mild transient and reversible thrombocytopenia, which appears not to be related to the amount of product infused, has been reported in some cases [1,6,7], suggesting that it may not be due to low levels of contaminating porcine vWF. The restoration of hemostasis with P-FVIII therapy is usually associated with increased measurable functional factor VIII levels in patient plasmas. Some patients, however, demonstrate a good hemostatic response to treatment with P-FVIII concentrate in the absence of increased factor VIII levels [8,9]. We investigated whether P-FVIII activates platelets and thus

induces a hemostatic effect due to an acquired "hypercoagulability" state separate from that attributable to increased circulating FVIII. The results indicate that, in contrast to other coagulation factor replacement products, P-FVIII activates platelets and generates platelet microparticles.

MATERIALS AND METHODS

Therapeutic Coagulation Products Used

Porcine factor VIII (HYATE:C, Porton Speywood Ltd, Wrexham, UK), intermediate purity FVIII (Hae-

Contract grant sponsor: Speywood Laboratories; Contract grant sponsor: Canadian Hemophilia Society.

*Correspondence to: Dr. John Freedman, Director, Transfusion Medicine, St. Michael's Hospital, 30 Bond Street, Toronto, Ontario, Canada, M5B 1W8.

Received for publication 21 March 1997; Accepted 10 September 1997

mate-P, Behringwerke AG, Marburg, Germany), a high purity monoclonal antibody-purified FVIII (Hemofil M, Baxter Corp., Glendale, CA), recombinant FVIII (rFVIII, Kogenate, Bayer Corp., Elkhart, IN), and FEIBA (Immuno AG, Vienna, Austria) were obtained from routine stocks. Recombinant factor VIIa (rFVIIa) was a gift from K. Lawday (Novo Nordisk Canada Inc., Mississauga, Ont.). FEIBA was used at a final concentration of 5 U/ml and rFVIIa at a final concentration of 1 KIU/ml, while the other products were used at final concentrations of 2 U/ml. These concentrations were selected to simulate plasma concentrations in a patient after therapy. Cryoprecipitate and cryosupernatant plasmas were obtained from Toronto Centre of the Canadian Red Cross Society Blood Services and were used at a 1:4 ratio v/v product to blood. "Aged" cryoprecipitate was prepared by maintaining freshly thawed cryoprecipitate at 4°C for 96 hr. Purified porcine vWF (126 µg/ml) was a gift from Dr. J.S. Pete Lollar, Emory University, Atlanta, GA.

Preparation and Staining of Platelets Using a Whole Blood Method

Using a 21-gauge needle and light tourniquet, after discarding the first 2 ml of blood, blood from normal healthy laboratory volunteers was drawn into citrated tubes (0.129 M citrate, Becton Dickinson, Rutherford, NJ). The citrated blood was diluted 1:2 with 0.35% bovine serum albumin/phosphate-buffered saline (BSA/PBS), pH 7.4, and supplemented with the synthetic peptide Gly-Pro-Arg-Pro (GPRP, Sigma Chemicals, St. Louis, MO) at a final concentration of 2.5 mM, to prevent thrombin-induced fibrin polymerization and platelet aggregation. As a positive control, human thrombin (Fibrindex, Ortho Diagnostics, Raritan, NJ) was added to the diluted blood sample to a final concentration of 1 U/ml and incubated at 37°C for 10 min. As negative controls (baseline), 0.35% BSA/PBS was added to the diluted blood. To test the effects of the therapeutic coagulation products on platelets, the diluted blood samples were incubated with each product for 30 min at 37°C. Following exposure to agonist, the samples were fixed with an equal volume of 1% (w/v) paraformaldehyde, pH 7.4, for 10 min at 22°C, as previously described [10]. The fixed samples were further diluted 1:5 with 0.35% BSA/PBS. Twenty-five microliters of fixed blood was incubated with saturating concentrations of monoclonal antibody in the dark for 30 min at 22°C [10]. Fluorescein isothiocyanate (FITC)-conjugated monoclonal antibodies to CD41 (GPIIb/IIIa), CD42 (GPIb), CD36 (GPIV; thrombospondin receptor), CD63 (lysosomal protein), and activated-GPIIb/IIIa (measuring the activation-induced conformationally changed CD41) were obtained from Immunotech Coulter, Westbrook, ME. Phycoerythrin (PE)-conjugated anti-CD42 and CD62P (P-selectin) were from Becton Dickinson, San Jose, CA. After incu-

bation, 1 ml of filtered FACSFlow fluid (Becton Dickinson) was added to each tube and the samples were acquired, within 4 hr after staining, using a FACScan flow cytometer (Becton Dickinson), equipped with a 15 mW argon ion laser. Gating on forward light scatter and fluorescence, GPIb- or GPIIb/IIIa-positive single platelets were identified, 10,000 events acquired on each sample and analyzed using LYSYS II software (Becton Dickinson). Quantitation of the proportion of cells expressing the marker or, using Simply Cellular beads (Flow Cytometry Standards, San Juan, PR), quantitation of the number of "molecules per cell" (antibody binding sites), was performed as described previously [10]. Mepacrine staining for platelet dense granules followed the method described by Gordon et al. [11].

Detection of Platelet Microparticles by Flow Cytometry

Microparticles were distinguished from intact normal platelets on the basis of the characteristic flow cytometric profile of forward light scatter vs. fluorescence with conjugated antiplatelet antibodies, as described by Abrams et al. [12]. The percentage of platelet microparticles was defined as that proportion of all GPIb- or GPIIb/IIIa-positive cells that were smaller than 0.8 µ; they showed reduced fluorescence compared to intact platelets, as shown in Figure 1.

Statistical Analysis

Since the basal level of the markers measured differs between different donors, results were "normalized" by relating each result to that of the baseline unstimulated resting platelets for that platelet donor, i.e., by calculating the ratio of post:pre exposure to agonist. One-way analysis of variance and paired Student's *t*-test were performed and *P* < 0.05 was regarded as significant.

RESULTS

Activation-Induced Translocation of Intracellular Molecules to Platelet Surface

Table I shows that following exposure of resting platelets to thrombin, there was an increase in platelets expressing surface CD62 (means ± SD, 3 ± 2% prestimulation vs. 96 ± 3% after thrombin; *P* < 0.0001), P-FVIII also increased platelets expressing CD62 (5-fold, *P* = 0.0118), but to a lesser extent (*P* < 0.001) than was observed with thrombin. Hemofil M, recombinant FVIII, recombinant FVIIa, Haemate P, and FEIBA did not increase the number of platelets expressing CD62. Similarly, P-FVIII increased the proportion of platelets expressing CD63 (5-fold), again to a lesser extent (*P* <




Fig. 1. Flow cytometric dot plot analysis of platelet microparticles, showing two populations of CD41 positive events; the larger cells (right) represent intact platelets, the smaller cells, (left) represent platelet microparticles. FSC = forward light scatter, reflecting the size of the cells; FL1 = fluorescence intensity of bound FITC-labeled anti-CD41.

TABLE I. Proportion of Platelets (mean \pm SD) Expressing CD62, CD63, and Activated GPIIb/IIIa, Following Incubation With Thrombin and Therapeutic Coagulation Products*

Agonist	N	CD62 (%)	CD63 (%)	Activated GPIIb/IIIa (%)
Unstimulated (baseline)	20	3.3 \pm 1.8	2.3 \pm 1.1	1.3 \pm 0.8
Thrombin	20	96 \pm 3	71 \pm 8	8.8 \pm 4.5
P-FVIII	20	15 \pm 13	11 \pm 6	8.6 \pm 9.1
Hemofil M	8	4.8 \pm 2.8	2.9 \pm 2.1	0.8 \pm 0.5
Haemate-P	8	1.6 \pm 1.0	1.7 \pm 0.4	1.3 \pm 0.9
rFVIII	8	3.1 \pm 1.2	1.8 \pm 0.6	0.6 \pm 0.2
FEIBA	8	1.9 \pm 1.1	0.7 \pm 0.3	3.0 \pm 1.2
rFVIIa	8	1.9 \pm 0.6	0.9 \pm 0.6	0.8 \pm 0.6
Cryosupernatant plasma	4	1.4 \pm 0.7	0.8 \pm 0.3	2.4 \pm 0.9
Cryoprecipitate, fresh	8	17 \pm 10	4.5 \pm 1.4	2.0 \pm 1.1
Cryoprecipitate, "aged"	8	6.0 \pm 3.7	2.7 \pm 0.5	2.1 \pm 1.4

*N = number of experiments.

0.0001) than was seen when thrombin was used as the agonist.

Activation-Induced Conformational Change in Normal Platelet Surface Glycoprotein

Few ($1 \pm 1\%$) unstimulated resting platelets showed expression of activated GPIIb/IIIa. As shown in Table I, exposure to thrombin in an increase in the proportion of platelets expressing activated GPIIb/IIIa. Porcine FVIII also increased activated GPIIb/IIIa expression (8-fold, $P = 0.002$), and, in contrast to CD62 and CD63, this was not significantly different from the values observed with thrombin ($P = 0.5954$). None of the other coagulation

products tested significantly increased platelet-activated GPIIb/IIIa expression.

Activation-Induced Modulation of Normal Platelet Surface Glycoproteins Expression

Expression of CD41, the fibrinogen receptor, is increased upon platelet activation [13]. Resting platelets expressed $49,812 \pm 5383$ (means \pm SD) CD41 molecules per cell ($n = 17$). As shown in Figure 2, thrombin markedly increased CD41 expression (a 1.74-fold increase, $P < 0.0001$). A significant increase (1.27-fold, $P < 0.0001$) in CD41 expression was also induced P-FVIII, but this was significantly lower than that seen with thrombin ($P < 0.0001$). None of the other coagulation products tested caused an increase in platelet surface CD41 expression.

Platelet surface expression of CD42, the vWF receptor, characteristically is reduced in this system following exposure to thrombin agonist [10,13]. The mean \pm SD CD42 on resting platelets was $22,345 \pm 4,878$ molecules/cell, $n = 17$. As shown in Figure 2, thrombin exposure reduced the expression of CD42 by a mean of 28% ($P < 0.0001$). In contrast, P-FVIII slightly increased the levels of CD42 expression (mean 9% increase, $P = 0.0014$). None of the other coagulation products significantly altered CD42 expression.

Platelet surface CD36 expression is also characteristically increased after platelet activation. The mean \pm SD CD42 on resting platelets was $22,345 \pm 4,878$ molecules/cell, $n = 17$. Figure 2 indicates that P-FVIII increased CD36 expression to the same extent as was seen with thrombin. Exposure of platelets to the other coagulation products did not increase CD36 expression.




Fig. 2. Ratio of platelet surface expression of CD41, CD42, and CD36 and microparticle generation following exposure to thrombin and to porcine factor VIII compared to levels on basal (unstimulated) resting platelets.

Analysis of Platelet Dense Granules

In contrast to the characteristic thrombin-induced reduction in platelet dense granules, none of the coagulation products tested affected platelet dense granule staining (data not shown).

Microparticle Formation

Platelet activation may be associated with generation of platelet fragments or microparticles. In the resting platelet samples, $4.0 \pm 0.8\%$ of all cells positive for the CD41 were identified as microparticles by their size. While exposure to the other coagulation products did not result in an increase in microparticle generation, as shown in Figure 2, exposure of platelets to either thrombin or P-FVIII significantly increased microparticles (2-fold increase with thrombin and 1.8-fold increase with PFVIII, $P = 0.0002$ and $P < 0.0001$, respectively). The extent of microparticle generation was not different between thrombin and P-FVIII ($P = 0.3293$).

Exposure of Platelets to Cryoprecipitate and Cryosupernatant Plasma

Exposure to cryosupernatant plasma did not result in platelet activation by any of the measurements employed. As shown in Table I, exposure of platelets to cryoprecipitate appeared to increase the number of platelets expressing CD62 (to the same extent [$P = 0.9$] as seen with P-FVIII), and this effect was attenuated with "aged" cryoprecipitate (a 2-fold increase over baseline, $P = 0.023$; $P = 0.001$ compared to fresh cryoprecipi-

tate). Cryoprecipitate did not significantly increase the number of platelets expressing CD63 or activated GPI-IbIIIa (Table I). While exposure of platelets to cryoprecipitate did not affect CD41, CD42, or CD36 platelet surface expression, the samples incubated with cryoprecipitate appeared to contain the greatest amount of microparticle formation (a 5.2-fold increase over unexposed platelets, $P < 0.0001$) and this effect was less marked with "aged" cryoprecipitate (3.3-fold increase, $P = 0.009$, not shown). However, subsequent examination of cryoprecipitate (both fresh and aged) alone, i.e., without incubation with platelets, showed that platelet microparticles were already present in the product, in an amount sufficient to explain the apparent increase observed when incubated with test platelets.

Exposure of Platelets to Porcine vWF

Platelets were also incubated under identical conditions with varying concentrations (serial dilutions from $0.0002 \mu\text{g/ml}$ to $50 \mu\text{g/ml}$) of porcine vWF. Limited studies ($n = 3$) showed that, with the exception of the slight increase in CD42 expression seen with P-FVIII, the porcine vWF yielded results similar to the P-FVIII preparation when used at a concentration of $1\text{--}2 \mu\text{g/ml}$. No effect on microparticle generation or platelet activation in any of the parameters measured was seen at lower concentrations and little or no effect was seen at higher concentrations other than a progressive blockade of surface CD42 sites with an increasing concentration of porcine vWF.

DISCUSSION

The induction of human platelet aggregation *in vitro* by P-FVIII was recognized in 1973 [14] and was shown to be due to porcine vWF present in the FVIII concentrates [3–5]. Porcine vWF can bind to platelet CD42 and activate the CD41 receptor on human platelets [4]; the binding of porcine vWF to CD42 in the absence of any inducers, such as ristocetin or botrocetin, results in platelet-platelet interaction, which leads to platelet aggregation. The thrombocytopenia observed with early preparations of P-FVIII was attributed to the platelet aggregation. Newer preparations of P-FVIII (HYATE:C) are more highly purified freeze-dried concentrates of porcine antihemophilic factor, prepared by ion-exchange chromatography with polyelectrolytes to separate and remove porcine vWF [15], and are generally associated with less significant thrombocytopenia post-infusion [6–8].

We examined platelet activation following incubation of normal human platelets with various therapeutic products employed in the management of patients with coagulation deficiencies. Incubation with monoclonal purified FVIII, recombinant FVIII, Haemate-P, FEIBA, recombinant FVIIa, and cryosupernatant plasma, did not result in platelet activation. Incubation with P-FVIII, in contrast, resulted in an increase in platelets expressing CD62, CD63, and activated GPIIb/IIIa, increased platelet surface expression of CD36 and CD41. These changes were consistent with those observed with thrombin-induced platelet activation. Fewer platelets expressed activated GPIIb/IIIa after exposure to thrombin and P-FVIII than those expressing CD62 and CD63 (Table I); this relates to the requirement for fibrinogen to demonstrate altered GPIIb/IIIa conformer, but in the test system employed the plasma is diluted 32-fold, reducing available fibrinogen. Although the same concentrations of thrombin, P-FVIII, and platelets were used throughout, with some markers of platelet activation the magnitude of the activation-induced changes with P-FVIII was lower than seen with thrombin (e.g., CD62, CD63, CD41), but for others (e.g., CD36 and activated GPIIb/IIIa) the magnitude of change was comparable. Thrombin caused a decrease in CD42 expression on the platelet surface, whereas P-FVIII caused a slight, but significant, increase. Although we have examined platelet activation in a variety of clinical disorders and under a variety of *in vitro* conditions, this is the only platelet agonist that caused an increase in platelet surface CD42 expression with platelet activation; the explanation for this observation remains to be determined. The differences in degree of platelet activation with some markers suggest that the mechanism of activation may be different between thrombin and P-FVIII. The observations with porcine vWF suggest that the platelet activation and microparticle generation induced by therapeutic concentrations of

P-FVIII (HYATE:C) are due to small amounts of residual porcine vWF in the preparation. Preliminary studies suggest that there is an optimal concentration of porcine vWF to achieve these effects. As discussed below, the effects of residual porcine vWF may be desirable.

The apparent platelet microparticle generation observed following incubation with cryoprecipitate seems to be due to platelet microparticles already present in the cryoprecipitate product; this is consistent with previous observations ([16] and personal communication, Georges Rivard, Montreal, 1996) and may be due to the processing, freezing, and thawing of these products.

Platelet activation has been associated with hypercoagulable and thrombotic states [17,18]. Platelet microparticles are formed when there is strong platelet activation. They contain platelet membrane CD41 and CD42, as well as membrane cytoskeletal proteins [19,20]. Microparticles derived from platelet activation are highly enriched in factor Va and Xa binding sites and exhibit prothrombinase activity [19,21]; they can also bind factors VIIIa and IXa [22,23]. In addition, platelet microparticles adhere to the subendothelium and promote platelet binding [24]; they provide a catalytic surface for production of both factor Xa and thrombin [19,20]. Infusion of microparticles shortens the bleeding time and the activated partial thromboplastin time in thrombocytopenic rabbits [25,26], and Warkentin et al. suggested that generation of procoagulant microparticles could explain the thrombotic complications in patients with heparin-induced thrombocytopenia [27]. Others have suggested that increased circulating platelet microparticles may increase the risk of thromboembolic events in patients with hemolytic uremic syndrome and thrombotic thrombocytopenic purpura [28], transient ischemic attacks, lacunar infarcts, and multi-infarct dementias [29,30].

Although the efficacy of P-FVIII (HYATE:C) generally correlates with the increment in circulating factor VIII achieved, a number of good or fair responses have been observed in the absence of a measurable factor VIII increment [8]. This somewhat surprising observation was confirmed by French investigators [9]. Whether the *in vitro* observations reported here occur *in vivo* is currently under investigation. Based on our findings to date, we hypothesize that the platelet activation and microparticle generation following infusion of P-FVIII may induce a hypercoagulable state that could explain the enhanced hemostatic response to P-FVIII observed in some hemophiliacs even in the absence of increased circulating factor VIII levels.

ACKNOWLEDGMENTS

This work was supported in part by a grant from Speywood Laboratories, Wrexham, UK and by a grant from the Canadian Hemophilia Society.

REFERENCES

- Gringeri A, Santagostino E, Tradati F, Giangrande PLF, Mannucci PM: Adverse effects of treatment with porcine factor VIII. *Thromb Haemost* 65:245, 1991.
- Lozier JN, Santagostino E, Kaspar CK, Teitel JM, Hay CR: Use of porcine factor VIII for surgical procedures in hemophilia A patients with inhibitors. *Semin Hematol* 30:10, 1993.
- Altieri DC, Capitanio AM, Mannucci PM: Von Willebrand factor contaminating porcine factor VIII (Hyate: C) causes platelet aggregation. *Br J Haematol* 63:703, 1986.
- Pareti FI, Mazzucato M, Bottini E, Mannucci PM: Interaction of porcine factor von Willebrand factor with the platelet glycoproteins Iband IIb/IIIa complex. *Br J Haematol* 82:81, 1992.
- Mazzucato M, De Marco L, Pradella P, Masotti A, Pareti FI. Porcine von Willebrand factor binding to human platelet GPIb induces transmembrane calcium influx. *Thromb Haemost* 76:655, 1996.
- Kernoff PBA, Thomas ND, Lilley PA, Matthews KB, Goldman E, Tuddenham EGD: Clinical experience with polyelectrolyte fractionated porcine factor VIII concentrate in the treatment of hemophiliacs with antibodies to Factor VIII. *Blood* 63:31, 1984.
- Brettler DB, Forberg AD, Levine PH, Aledort LM, Hilgartner MW, Kasper CK, Lusher JM, McMillan C, Roberts H, The Cooperating Investigators: The use of porcine factor VIII concentrate (HYATE:C) in the treatment of patients with inhibitor antibodies to factor VIII. *Arch Int Med* 1249:1381, 1989.
- Hay CRM, Lozier JN, Lee CA, Lafan M, Tradati H, Santagostino E, Ciavarella N, Schiavoni M, Fukui H, Yoshioka A, Teitel J, Mannucci PM, Kasper CK: Porcine factor VIII therapy in patients with congenital hemophilia and inhibitors: efficacy, patient selection, and side effects. *Semin Hematol* 31(Suppl 4):20, 1994.
- Negrier C: The use of porcine factor VIII in France. *Semin Hematol* 31(Suppl 4):29, 1994.
- Kühne T, Hornstein A, Semple J, Chang W, Blanchette V, Freedman J. Flow cytometric evaluation of platelet activation in blood collected into EDTA vs. Diatube-H, a sodium citrate solution supplemented with theophylline, adenosine, and dipyridamole. *Am J Hematol* 50:40, 1995.
- Gordon N, Thom J, Cole C, Baker R: Rapid detection of hereditary and acquired platelet storage pool deficiency by flow cytometry. *Br J Haematol* 89:117, 1995.
- Abrams CS, Ellison N, Budzynski AZ, Shattil SJ: Direct detection of activated platelets and platelet-derived microparticles in humans. *Blood* 75:128, 1990.
- Michelson AD, Barnard MR: Thrombin-induced changes in platelet membrane glycoproteins Ib, IX, and IIb-IIIa complex. *Blood* 70:1673, 1987.
- Forbes CD, Prentice CRM: Aggregation of human platelets by purified porcine and bovine antihemophilic factor. *Nature: New Biology* 241:149, 1973.
- Middleton S: Polyelectrolytes and preparation of Factor VIII:C. In Forbes CD, Lowe GD (eds): "Unresolved Problems in Hemophilia." Lancaster: MTP Press, 1982, pp 109-120.
- George PN, Pickett EB, Heinz R: Platelet membrane microparticles in blood bank fresh frozen plasma and cryoprecipitate. *Blood* 68:307, 1986.
- Michelson AD: Flow cytometry: A clinical test of platelet function. *Blood* 87:4925, 1996.
- Janes SJ, Wilson DJ, Chronos N, Goodall AH: Evaluation of whole blood flow cytometric detection of platelet bound fibrinogen on normal subjects and patients with activated platelets. *Thromb Haemost* 70:659, 1993.
- Sims PJ, Wiedmer T, Weiss HJ, Shattil SJ: Assembly of the platelet prothrombinase complex is linked to vesiculation of the platelet plasma membrane. *J Biol Chem* 264:17049, 1989.
- Fox JB, Austin CD, Reynolds CC, Steffen PK: Evidence that agonist-induced activation of calpain causes the shedding of procoagulant-containing microparticles from the membrane of aggregating platelets. *J Biol Chem* 266:13289, 1991.
- Wiedmer T, Shattil SJ, Cunningham M, Sims PJ. Role of calcium and calpain in complement-induced vesiculation of the platelet plasma membrane and in the exposure of the platelet Factor Va receptor. *Biochemistry* 29:623, 1988.
- Gilbert JE, Sims PJ, Wiedmer T, Furie B, Furie BC, Shattil SJ: Platelet-derived microparticles express high affinity receptors for Factor VIII. *J Biol Chem* 266:17261, 1991.
- Hoffman M, Munove DM, Roberts HR: Coagulation factor IXa binding to activated platelets and platelet derived microparticles: A flow cytometry study. *Thromb Haemost* 68:74, 1992.
- Owens MR, Holme S, Cardinali S: Platelet microparticles adhere to subendothelium and promote adhesion of platelets. *Thromb Res* 66:2475, 1992.
- McGill M, Fugman DA, Vittorio N, Darrow C: Platelet membrane microparticles reduced microvascular bleeding times in thrombocytopenic rabbits. *Lab Clin Med* 109:127, 1987.
- Howard MA, Coghlan M, David R, Pfueller SL: Coagulation activities of plasma microparticles. *Thromb Res* 50:145, 1988.
- Warkentin TE, Hayward CPM, Boshkov L, Santos AV, Sheppard JA, Bode AP, Kelton JG: Sera from patients with heparin-induced thrombocytopenia generate platelet-derived microparticles with procoagulant activity: An explanation for the thrombotic complications of heparin-induced thrombocytopenia. *Blood* 84:3691, 1994.
- Galli M, Grassi A, Barbui T: Platelet-derived microparticles in thrombotic thrombocytopenic purpura and hemolytic uremic syndrome. *Thromb Haemost* 75:427, 1996.
- Lee YJ, Jy W, Horstman LL, Janania J, Reyes Y, Kelley RE, Ahn YS: Elevated platelet microparticles in transient ischemic attacks, lacunar infarcts, and multiinfarct dementia. *Thromb Res* 72:295, 1993.
- Jy W, Horstman LL, Arce M, Ahn YS: Clinical significance of platelet microparticles in autoimmune thrombocytopenias. *J Lab Clin Med* 119:334, 1992.